

Southeast Michigan Regional Protocol

Genesee, HEMS (Wayne), Lapeer, Macomb, Oakland, and Washtenaw/Livingston MCA's

Pediatric Altered Level Of Consciousness

Pre-Radio

MFR/EMT/SPECIALIST/PARAMEDIC

1. Follow General Pre-Hospital Care Protocol.

SPECIALIST/PARAMEDIC

2. Start an IV NS KVO.
3. If hypotensive, bolus with 20 ml/kg of NS.

PARAMEDIC

4. Check glucose level. If glucose is less than 80 mg/dl, administer Dextrose 2 ml/kg IV.
 - a. Dextrose 12.5% for children under 1 yr.
 - b. Dextrose 25% for children between 1 yr and 12 yrs.

PARAMEDIC

5. Administer Naloxone (Narcan) 0.5 mg for ages 0-2 yrs., 1 mg for ages 2-6 yrs., and 2 mg IV/IM for patients more than 6 yrs.

Post-Radio

PARAMEDIC

6. Repeat Dextrose as indicated.
7. Repeat Naloxone (Narcan) as indicated.

NOTE: To obtain approximately 12.5% Dextrose mixture draw 37.5 ml out of one amp of D50 and discard, then add 37.5 ml of NS; administer as indicated above.

To obtain approximately 25% Dextrose mixture, draw 25 ml out of one amp of D50 and discard, then add 25 ml of NS; administer as indicated above.