

HIV/AIDS DATA

WASHTENAW COUNTY RESIDENTS

Newly diagnosed HIV cases:

	2002	2003	2004	2005	2006	2007	2008
Total HIV cases	20	28	31	27	29	28	30
Male	17	22	24	24	24	22	25
Female	3	6	7	3	5	6	5
Black	10	12	17	10	14	16	16
White	9	13	11	15	13	11	13
Age 15 to 29 yr	3	11	13	9	4	16	16
Anonymous cases (included in total HIV)	6	5	6	3	1	5	5

Highest zip codes for new HIV cases:

	2002	2003	2004	2005	2006	2007	2008
48197	30%	14%	23%	52%	21%	31%	48%
49198	15%	27%	38%	15%	34%	27%	24%

Reported risk factors in new HIV cases:

	2002	2003	2004	2005	2006	2007	2008
MSM¹	65%	50%	52%	70%	64%	57%	70%
Bisexual male	10%	11%	16%	19%	21%	21%	13%
IV Drug Use	10%	7%	3%	7%	0%	0%	3%
Hetero, high risk*	5%	4%	13%	7%	7%	4%	3%
Hetero, low risk**	10%	21%	29%	15%	29%	32%	20%
Hx. Blood products	0%	5%	0%	0%	0%	0%	0%

¹MSM = Men who have sex with men (includes bisexual males)

*Hetero, high risk = Heterosexual sex with IDU, known HIV+ partner, bisexual male, sex worker etc.

**Hetero, low risk = Heterosexual sex with no other risk identified

Note: Risk percentages are calculated in relation to the total number of cases. Risk percentages may add up to more or less than 100% due to some individuals having more than one risk and others having none identified.

Note: HIV+ data include anonymous and confidential cases, and exclude prison cases.

New AIDS cases:

	2002	2003	2004	2005	2006	2007	2008
Total new cases	16	6	12	19	14	11	8
Male	14	5	10	15	12	9	4
Female	2	1	2	4	2	2	4
Black	7	4	4	10	6	8	6
White	8	2	7	7	6	1	2
Co-diagnosis***	45%	11%	6%	22%	21%	7%	13%

*** Co-diagnosis = Proportion of HIV cases diagnosed with AIDS within 3 months of initial HIV diagnosis.